

KHSRP polyclonal antibody

Catalog: BS90767

Host: Rabbit

Reactivity:

ity: Human, Mouse, Rat

BackGround:

Binds to the dendritic targeting element and may play a role in mRNA trafficking (By similarity). Part of a ternary complex that binds to the downstream control sequence (DCS) of the pre-mRNA. Mediates exon inclusion in transcripts that are subject to tissue-specific alternative splicing. May interact with single-stranded DNA from the far-upstream element (FUSE). May activate gene expression. Also involved in degradation of inherently unstable mRNAs that contain AU-rich elements (AREs) in their 3'-UTR, possibly by recruiting degradation machinery to ARE-containing mRNAs.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

73 kDa

Swiss-Prot:

Q92945(Human) Q3U0V1(Mouse) Q99PF5(Rat)

Purification&Purity:

ProA affinity purified

Applications:

WB:1:500-1:2,000 ICC/IF:1:50-1:200 IHC:1:50-1:200 IP:1:10-1:50

Storage&Stability:


Store at +4 $^{\circ}$ C after thawing. Aliquot store at -20 $^{\circ}$ C. Avoid repeated freeze / thaw cycles.

Specificity:

KHSRP polyclonal antibody detects endogenous levels of

KHSRP protein.

DATA:




Western blot analysis of KHSRP on different lysates using anti-KHSRP antibody at 1/1,000 dilution.

Positive control:

Lane 1: 293

Lane 2: mouse brain

Lane 3: rat brain



ICC staining KHSRP in 293T cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151