

MRP2 polyclonal antibody

Catalog: BS90880

Host: Rabbit

Reactivity: Human

BackGround:

Multi-drug resistance protein 2 (MRP2), also known as ABCC2, is an ATP binding cassette (ABC) transporter responsible for biliary excretion of xenobiotics, endobiotics, and their metabolites. Deficiency in ABCC2 results in the clinical disorder Dubin-Johnson syndrome. MRP2 is found to be expressed in a variety of human cancers, and is associated with resistance of tumor cells to various anticancer drugs including cisplatin. The predicted molecular weight of MRP2 is 174 kDa, while mature MRP2 usually has a slower migration around 190-250 kDa due to the glycosylation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

174 kDa

Swiss-Prot:

Q92887(Human)

Purification&Purity:

ProA affinity purified

Applications:

WB:1:500-1:1,000

ICC:1:50-1:200

IHC:1:50-1:200

FC:1:50-1:100


Storage&Stability:

Store at +4 °C after thawing. Aliquot store at -20 °C or -80 °C. Avoid repeated freeze / thaw cycles.


Specificity:

MRP2 polyclonal antibody detects endogenous levels of MRP2 protein.

DATA:



ICC staining MRP2 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining MRP2 in 293T cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151