

GADD 45 γ (S123) polyclonal antibody

Catalog: BS9140

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Cell cycle progression is subject to arrest at G1 and G2 checkpoints in response to DNA damage, presumably to allow time for DNA repair prior to entry into S and M phase, respectively. The p53 tumor suppressor is required for one such G1 checkpoint and functions to upregulate expression of GADD 45 and p21. GADD 45 binds both Cdks and PCNA, a protein involved in DNA replication and repair. GADD 45 stimulates DNA excision repair in vitro and inhibits entry of cells into S phase. Thus, it has been suggested that GADD 45 may serve as a link between the p53-dependent cell cycle checkpoint and DNA repair. GADD 45-like proteins, GADD 45 β and GADD 45 γ , have been shown to be induced by environmental stresses. GADD 45 β and GADD 45 γ are thought to induce p38/JNK activation via MEKK4 activation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 17 kDa

Swiss-Prot:

O95257

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

GADD 45 γ (S123) polyclonal antibody detects endogenous levels of GADD 45 γ protein.

DATA:



Immunohistochemistry (IHC) analyzes of GADD 45 γ (S123) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151