

Histone H3 (mono+di+methyl K79) polyclonal antibody

Catalog: BS94075

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

Eukaryotic histones are basic and water soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed super-helical turn sequentially to form chromosomal fiber. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form the octamer; formed of two H2A-H2B dimers and two H3-H4 dimers, forming two nearly symmetrical halves by tertiary structure. Over 80% of nucleosomes contain the linker Histone H1, derived from an intronless gene, that interacts with linker DNA between nucleosomes and mediates compaction into higher order chromatin. Histones are subject to posttranslational modification by enzymes primarily on their N-terminal tails, but also in their globular domains. Such modifications include methylation, citrullination, acetylation, phosphorylation, sumoylation, ubiquitination and ADP-ribosylation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

15 kDa

Swiss-Prot:

P68431(Human) P84243(Human) Q16695(Human)
Q6NXT2(Human) Q71DI3(Human) P68433(Mouse)
Q6LED0(Rat)

Purification&Purity:

ProA affinity purified

Applications:

WB:1:1,000-1:2,000

IHC:1:50-1:200


Storage&Stability:

Store at +4 °C after thawing. Aliquot store at -20 °C or -80 °C. Avoid repeated freeze / thaw cycles.


Specificity:

Histone H3 (mono+di+methyl K79) polyclonal antibody detects endogenous levels of Histone H3 protein only when mono/di-methylated at K79.

DATA:



Western blot analysis of Histone H3(mono+di+tri methyl K79) on different lysates using anti-Histone H3(mono+di+tri methyl K79) antibody at 1/1,000 dilution. Positive control: Lane 1: Mouse testis Lane 2: CRC



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Histone H3(mono+di+tri methyl K79) antibody. Counter stained with hematoxylin.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151