

polyclonal antibody Smad5

Catalog:	BZ16569	Host:	Rabbit	Reactivity:	Human,Rat
BackGrou	nd:			FC: 1/50-1/100	

Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. Smad5 is a receptor-regulated Smad (R-Smad). Smad5 is required for normal development of the cardiovascular system in vivo; lack of the Smad5 gene results in apoptosis of cardiac myocytes. 3 Upregulation of Smad5 has been reported to mediate apoptosis of gastric epithelial cells induced by Helicobacter pylori infection. Tissue specificity: Ubiquitous.

Product:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Molecular Weight:

Calculated MW: 52 kDa; Observed MW: 52 kDa

Swiss-Prot:

Q99717

Purification&Purity:

Affinity Purified

Applications:


WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 **Storage&Stability:**

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Isotype:

IgG

DATA:



Western blot analysis of Smad5 in HEK293 lysates using Smad5 anti-

body.

Note:

For research use only, not for use in diagnostic procedure.