

V5-Tag Mouse monoclonal antibody

Catal	log:	MB2020

Host: Mouse

Reactivity: All

BackGround:

The V5 epitope tag is derived from a small epitope (Pk) present on the P and V proteins of the paramyxovirus of simian virus 5 (SV5). The V5 tag is usually used with all 14 amino acids, although it has also been used with a shorter 9 amino acid sequence.

Product:

Mouse IgG1, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

N/A

Swiss-Prot:

N/A

Purification&Purity:

Affinity-chromatography

Applications:

WB: 1:5000~1:10000

IF: 1:200


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

V5-tag mouse monoclonal antibody detects endogenous levels of internal, C-terminal, and N-terminal V5- tagged proteins.

DATA:



All lanes : V5-tag mouse monoclonal antibody A: HEK 293 cell lysate transfected without V5-tag protein B: HEK 293 cell lysate transfected with V5 -tag protein C: IP (HEK 293 cell without transfected + anti- V5 mAb 4°C overnight) D: IP (PBS + anti- V5

Note:

For research use only, not for use in diagnostic procedure.