

ALDH4A1 monoclonal antibody

Catalog: MB9048

Host: Mouse

Reactivity: Human, Rat

BackGround:

Aldehyde dehydrogenases (ALDHs) mediate NADP⁺-dependent oxidation of aldehydes into acids during detoxification of alcohol-derived acetaldehyde, lipid peroxidation and metabolism of corticosteroids, biogenic amines and neurotransmitters. ALDH4A1 (aldehyde dehydrogenase 4 family member A1), also known as P5CD (Delta-1-pyrroline-5-carboxylate dehydrogenase), P5CDh, P5CDhL, P5CDhS or ALDH4, is a major enzyme involved in the proline degradation pathway. Localizing to the mitochondrial matrix, ALDH4A1 catalyzes the conversion of Delta-1-pyrroline-5-carboxylate (P5C) to glutamate. A mutation in the gene encoding ALDH4A1 results in HPII (hyperprolinemia type II), a disease characterized by an excess of P5C and proline that is associated with mental retardation and seizures.

Product:

Mouse IgG1, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

62 kDa

Swiss-Prot:

P30038(Human) P0C2X9(Rat)

Purification&Purity:

ProG affinity purified.

Applications:

WB:1:1,000-1:2,000

IHC:1:50-1:200

Storage&Stability:


Store at +4 °C after thawing. Aliquot store at -20 °C. Avoid

repeated freeze / thaw cycles.


Specificity:

ALDH4A1 monoclonal antibody detects endogenous levels of ALDH4A1 protein.

DATA:



Western blot analysis of ALDH4A1 on SK-Br-3 (1) and HepG2 (2) cell lysate using anti-ALDH4A1 antibody at 1/1,000 dilution.



Immunohistochemical analysis of paraffin-embedded human prostate tissue using anti-ALDH4A1 antibody. Counter stained with hematoxylin.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151