

PLGF monoclonal antibody

Catalog: MB9435

Host: Mouse

Reactivity: Human, Mouse, Rat

BackGround:

The onset of angiogenesis is believed to be an early event in tumorigenesis and may facilitate tumor progression and metastasis. Several growth factors with angiogenic activity have been described. These include fibroblast growth factor (FGF), platelet derived growth factor (PDGF), vascular endothelial growth factor (VEGF) and placenta growth factor (PIGF). Like VEGF, several PIGF variants have been shown to arise from alternative mRNA splicings. Evidence has suggested VEGF to be an obligatory component in PIGF signaling. While VEGF homodimers and VEGF/PIGF heterodimers function as potent mediators of mitogenic and chemotactic responses in endothelial cells, PIGF homodimers are effectual only at extremely high concentrations. Indeed, many of the physiological effects attributed to VEGF may actually be a result of VEGF/PIGF. VEGF and PIGF share a common receptor, Flt-1, and may also activate Flk-1/KDR.

Product:

Mouse IgG1, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

25 kDa

Swiss-Prot:

P49763(Human) P49764(Mouse) Q63434(Rat)

Purification&Purity:

ProG affinity purified.

Applications:

WB:1:500

ICC:1:50-1:100

IHC:1:50-1:200


Storage&Stability:

Store at +4 °C after thawing. Aliquot store at -20 °C. Avoid repeated freeze / thaw cycles.


Specificity:

PLGF monoclonal antibody detects endogenous levels of PLGF protein.

DATA:



Western blot analysis of PLGF on JAR using anti-PLGF antibody at 1/500 dilution.



ICC staining PLGF (green) in HepG2 cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151